

Study Confirms Link Between COVID-19 Vaccination and Menstrual Cycle Length

COVID-19 vaccination was associated with a temporary average increase in menstrual cycle length of less than one day.

October 4, 2022 By National Institutes of Health

Study confirms link between COVID-19 vaccination and temporary increase in menstrual cycle length

A large international study has confirmed the findings of a [previous U.S. study](#) that linked COVID-19 vaccination with an average increase in menstrual cycle length of less than one day. The increase was not associated with any change in the number of days of menses (days of bleeding).

Funded by the National Institutes of Health, the new study included data from nearly 20,000 people from Canada, the United Kingdom, the United States, Europe and other parts of the world who received any of nine different vaccines. For most study participants, the increase resolved in the cycle following vaccination.

The study's principal investigator was Alison Edelman, MD, MPH, of Oregon Health & Science University in Portland. [It appears in BMJ Medicine.](#)

"These findings provide additional information for counseling women on what to expect after vaccination," said Diana Bianchi, MD, director of NIH's Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD). "Changes following vaccination appear to be small, within the normal range of variation, and temporary."

NICHD and NIH's Office of Research on Women's Health funded the study, which was part of \$1.67 million awarded to five institutions to explore potential links between COVID-19 vaccination and menstrual changes.

A change in cycle length of less than eight days is considered within the normal range of variation. Although small menstrual changes may not be meaningful to health care professionals and researchers, the study authors wrote, perceived changes in a bodily function linked to fertility may be alarming to those experiencing it and could contribute to vaccine hesitancy.

Researchers analyzed de-identified data from the fertility tracking app, Natural Cycles. Users of the app provided it information on their temperature and their menstrual cycles. They also had the option to consent to the use of their de-identified data for research.

Worldwide rollout of COVID-19 vaccines allowed the study authors to expand on their original study of people in the United States. The study included data from participants throughout the world, but most were from the United Kingdom (32%), the United States and Canada (29%) and Europe (34%). In addition to the messenger RNA COVID-19 vaccines (Pfizer-BioNTech and Moderna), participants received COVID-19 vaccines made from engineered viruses (AstraZeneca, Covishield, Janssen/Johnson & Johnson and Sputnik), and inactivated viruses (Covaxin, Sinopharm, and Sinovac).

A total of 19,622 individuals participated. Of these, 14,936 were vaccinated and 4,686 were not. The researchers analyzed data on at least three consecutive cycles before vaccination and at least one cycle after. Data from at least four consecutive cycles were analyzed over a similar time interval for unvaccinated participants.

On average, vaccinated people experienced an increase of less than one day in each cycle in which they were vaccinated: a .71 day increase after the first dose and a .56 day increase after the second dose. Participants who received both doses in a single cycle had a 3.91 day increase in cycle length. After vaccination, cycle length had increased by only .02 days for individuals who received one dose per cycle, and .85 days for individuals who received two doses in one cycle, compared to participants who were not vaccinated. Changes in cycle length did not differ according to the type of vaccine received.

Of the total, 1,342 participants experienced a change in cycle length of eight or more days, comprising 6.2% of vaccinated individuals and 5.0% of unvaccinated individuals. Women who were younger and who had longer cycle length before vaccination were more likely to experience the increase.

The authors called for future studies on other aspects of vaccination-linked changes to menstrual cycles, such as unexpected vaginal bleeding and menstrual flow and pain. They also called for studies to determine the physical reasons why such changes might occur.

This [news release](#) was originally published by the National Institutes of Health on September 27, 2022.